

1 Semicircular canal size and locomotion in colobine monkeys: a cautionary tale

2

3 Todd C. Rae¹, Paul Martin Johnson¹, Wataru Yano² & Eishi Hirasaki³.

4

5 ¹ Centre for Research in Evolutionary, Social and Inter-Disciplinary Anthropology, University of
6 Roehampton, UNITED KINGDOM

7 ² Dept of Oral Anatomy, Asahi University School of Dentistry, JAPAN

8 ³ Primate Research Institute, Kyoto University, Inuyama, Aichi 484-8506, JAPAN

9

10 Running head: Semicircular canal size in colobines

11

12 Keywords: bony labyrinth, inner ear morphology, leaping, balance, *Colobus*, *Presbytis*

13

14 Word count: 3960

15

16 Address for correspondence:

17 Todd C. Rae

18 Centre for Research in Evolutionary, Social and InterDisciplinary Anthropology

19 Department of Life Sciences

20 University of Roehampton

21 Holybourne Avenue

22 London SW15 4JD

23 UNITED KINGDOM

24

25 email: t.rae@roehampton.ac.uk

26 Tel: +44 (0)20 8392 3726

27 Fax: +44 (0)20 8392 3610

28

29 ABSTRACT

30

31 The semicircular canals of the inner ear are the organ of balance, tracking head rotation during
32 movement and facilitating stabilisation of vision. Morphological characteristics of the canals are
33 correlated with agility scores related to locomotion. To date, however, the relationship between
34 canal morphology and specific locomotor behaviours, such as leaping, is unclear. Knowledge of
35 such a relationship could strengthen the inferences of locomotion of extinct taxa. To test this, crania
36 of two sets of closely related primate species (*Presbytis melalophos* and *P. potenziani*; *Colobus*
37 *guereza* and *C. polykomos*) that differ in the percentage of leaping in their locomotor repertoire
38 were examined using microscopic computed tomography (μ CT). Three-dimensional virtual models
39 of the bony labyrinth were derived and the radius of curvature of each of the three canals was
40 evaluated relative to cranial size. The findings are contradictory; one leaping form (*P. melalophos*)
41 differs from its congener in possessing significantly larger lateral canals, a pattern seen in previous
42 studies of primates, while the other leaper (*C. guereza*) has significantly smaller posterior canals
43 than its close relative. These results undermine efforts to determine specific locomotor behaviours
44 from the bony labyrinth of extinct primates.

45

46

47

48 INTRODUCTION

49

50 Locomotor studies of primates focus primarily on the postcranial evidence (e.g., Su and Jablonski,
51 2009). Hard and soft tissue characteristics generally associated with habitual locomotor modes are
52 found in the limbs (Fleagle, 1976, 1977), although non-locomotor activity also affects the
53 postcrania (Dunham et al., 2015). More recently, however, the advent of modern imaging
54 techniques has allowed the examination of the internal portions of the cranium, including the
55 labyrinth of the temporal bone, which includes the organ of balance (Spoor & Zonneveld, 1998).
56 Exploration of the link between labyrinth morphology and motion has yielded substantial results,
57 both within primates and among mammals more generally (Spoor et al., 2007; Welker et al., 2009;
58 Billet et al., 2012; Pfaff et al., 2015).

59

60 The inner ear (Fig. 1) is comprised of numerous connected cavities known as the osseous (or bony)
61 labyrinth in mammals (Gray, 1908; Spoor & Zonneveld, 1995, 1998; Schmelzle et al. 2007; Welker
62 et al., 2009; Billet et al., 2012; Gunz et al., 2012; Pfaff et al., 2015) and other vertebrates (Boistel et
63 al., 2011). The bony labyrinth consists of the cochlea, vestibule and semicircular canals
64 (Sinnatamby, 2006), in which are contained the components of the membranous labyrinth: cochlea
65 duct, the sac-like utricle and saccule, and the semicircular ducts, respectively (Gray, 1908;
66 Sinnatamby, 2006; Walker et al., 2008). The cochlea houses the hearing system, while the vestibular
67 system (vestibule and semicircular canals) forms the organ of balance. These systems reside within
68 the otic capsule of the petrosal portion of the temporal bone (Lieberman, 2011).

69

70 Figure 1 about here

71

72 The membranous labyrinth is smaller in diameter (lumen) than the enclosing osseous labyrinth, and
73 is separated from the bony wall by a supporting fluid known as perilymph (Ekdale, 2013). The

74 membranous labyrinth contains fluid known as endolymph (Spoor, 2003; Sinnatamby, 2006), a
75 heavy gelatinous fluid (Gibson, 1968) that responds to rotational acceleration of the head with a
76 corresponding deflection, which in turn moves the hair-like cilia inside the semicircular ducts
77 proportionally to the force of the turn; the corresponding signal is transmitted to the brain via the
78 vestibulocochlear nerve (Gibson, 1968; Mather, 2009). The semicircular canals (and the ducts
79 within them) are orientated in roughly orthogonal planes; thus, the brain interprets rotary head
80 movement through a combination of input in three different axes: pitch, roll and yaw (Walker et al.,
81 2008). It is the membranous labyrinth that determines the crucial factors for balance sensitivity
82 (e.g., size of the semicircular ducts), but as these variables are difficult or impossible to determine
83 after death, characteristics of the bony canals alone are used in studies of the relationship between
84 labyrinth morphology and locomotion (Spoor, 2003).

85

86 Figure 2 about here

87

88 At a relatively wide scale, placental mammal species that move in ways that include quicker and
89 more active actions require changes in the semicircular canals of the inner ear to cope with the more
90 extensive head movement these taxa experience (Spoor et al. 2007; Ryan et al., 2012). This
91 mammalian pattern has been proposed for primates, as well; for example, forms that habitually leap
92 (e.g., galagos, tarsiers) are reported to have relatively large semicircular canals, especially the
93 lateral (Spoor et al. 2007). For this finding to be operationalised with respect to fossil taxa,
94 however, it is necessary to demonstrate that similar and/or closely related taxa that differ in
95 locomotor pattern also differ significantly in bony labyrinth morphology, particularly as some extant
96 primates have been shown to deviate from expectation (Spoor et al., 2007). The hypothesis that
97 *Colobus* possesses larger canals than *Procolobus* because the former leaps more frequently (Walker
98 et al., 2008) is taken as the starting point of the present investigation.

99

100 To test the prediction that closely related colobines differ in their semicircular canal morphology
101 due to differences in the percentage of travel time spent leaping, an important part of the typical
102 colobine locomotor repertoire (Davison, 1982), two members of the genus *Colobus* (*C. polykomos*,
103 *C. guereza*) and the genus *Presbytis* (*P. melalophos* and *P. potenziani*) that differ in the percentage
104 of leaping in their overall locomotor repertoire (Table 1) are examined here. To support the
105 hypothesis that larger canals are associated with leaping, it is expected that both species pairs will
106 show the same relationship (i.e., the leaping form in each genus will have larger canals than its
107 congener).

108
109 All four of the taxa examined here are medium sized (6-13kg) arboreal monkeys with a high
110 percentage of leaves in the diet (Rowe, 1996; Fleagle, 1999). *P. melalophos* and *P. potenziani* live
111 on the SE Asian islands of Sumatra and Mentawai, respectively, although *P. melalophos* is also
112 found on the mainland of the Malay Peninsula (Fleagle, 1999). The difference in their degree of
113 leaping behaviour is not obviously related to their habitats, as both species inhabit primary and
114 secondary forests (Rowe, 1996). *C. polykomos* and *C. guereza* are larger and more dimorphic than
115 the Asian forms, and live in the western and central/eastern regions of the central African rainforest,
116 respectively (Fleagle, 1999). The percentage of travel time spent leaping differs between them
117 perhaps because *C. polykomos* lives primarily in continuous, primary forest, while *C. guereza*
118 frequents more secondary forest and wooded grassland (Rowe, 1996), where leaping between
119 supports is more likely to be necessary.

120
121 Phylogenetically, *Colobus* and *Presbytis* are both members of the Colobinae, but belong to separate
122 branches of the subfamily; *Presbytis* is more closely related to the 'odd nosed' colobines (*Nasalis*,
123 *Rhinopithecus*, *Pygathrix*), while *Colobus* is part of a monophyletic African clade with *Pliocolobus*
124 (Wang et al., 2012). In both cases, the species examined here are very closely related to their
125 congeners, sharing a common ancestor with only one other species (*P. comata* and *C. vellerosus*,

126 respectively), aside from each other (Ting, 2008; Meyer et al., 2011).

127

128 The decision to examine congeners was taken to diminish the effect of extraneous variables on the
129 ensuing comparisons; this can be useful for detecting adaptive responses (Crews, 1997). Species
130 pairs form natural experiments from which behaviours can be linked to morphology (Fleagle, 1976,
131 1977; Katz and Harris-Warrick, 1999). For example, comparing closely related taxa has the effect
132 of reducing the influence of phylogeny, so that those differences of interest (in this case, frequency
133 of leaping) can be examined in relative isolation (Birkhead et al., 1992). Using two different
134 species pairs eliminates many of the inherent problems (Garland and Adolf, 1994) with standard
135 two-species comparisons.

136

137 Table 1 about here

138

139 Materials and methods

140

141 A mixed-sex sample of wild-shot, adult dry crania of *P. melalophus* (n = 5) and *P. potenziani* (n = 4)
142 from the collection of the Primate Research Institute of Kyoto University, Inuyama, Japan, and of
143 *C. polykomos* (n = 6) and *C. guereza* (n = 5) from the Natural History Museum, London, U.K., was
144 examined. *Colobus* individuals were scanned at Hull York Medical School using a X-TEK
145 HMX160 high-resolution computed tomography scanner (X-Tek Systems, Tring, U.K.), with no
146 additional filters applied. All scans were taken at 75 Kv and 30 μ A. Voxel size was specimen-
147 dependent, but was ≤ 0.04 mm in all cases. *Presbytis* scan sets were obtained at the Asahi University
148 School of Dentistry, Japan, using a Scan-Xmate-RB090SS (Comscan Tecno, Sagamihara, Japan).
149 All scans were acquired at 80-90 Kv and 90-100 μ A, with a voxel size ≤ 0.04 mm.

150

151 Semicircular canal parameters were determined using a modified version (Schmelzle et al. 2007) of

152 the methods outlined in Spoor et al. (2007). Virtual 3D models of the labyrinth (Fig. 2) were
153 constructed via manual segmentation in Avizo ver. 8 (Visualization Sciences Group, Burlington,
154 MA). Although other parameters of vestibular morphology have been investigated (Pfaff et al.,
155 2015; Grohé et al., 2016), the present study focuses on the radius of curvature (R) as a measure of
156 canal size (Schmelzle et al., 2007). R is derived from internal height and width measurements:
157 height is measured from the protrusion of the ampulla at the lowest point of the canal to its
158 maximum possible extent on the canal opposite, width was then determined as the maximum length
159 line intersecting the height line at 90° (Fig. 3). R is calculated as $0.5 \times ((\text{Height} + \text{Width}) \div 2)$.

160

161 Figure 3 about here

162

163 To scale for size, values for each canal were derived by dividing R by the cranial body mass
164 estimate prosthion–inion cranial length (Martin and Ross, 2005); previous work (Johnson, 2012) on
165 *Colobus* shows that similar results obtain using other canal variables and scalars. The resulting
166 data were analysed for normality, then by independent samples tests in IBM SPSS ver. 21 with $\alpha =$
167 0.05 for significance.

168

169 Results

170

171 Examples of the virtual reconstructions of the canals are shown in Figure 4. The mean values of R
172 and cranial length are given in table 2. The 95% confidence intervals of the regression slopes of R
173 against size included isometry except in the case of the anterior canal, where the lower confidence
174 interval approached isometry. No samples of R for which $n > 5$ differed significantly from normality
175 (the small sample of *P. potenziani* precluded analysis).

176 Neither species pair differed significantly in size-corrected R for anterior canals (Fig. 5). In

177 *Presbytis*, however, the species that leaps more frequently (*P. melalophos*) shows significantly
178 larger lateral canals; $t = 2.867$, $df = 7$, Sig. (2-tailed) = 0.024. The results for *Colobus*, however,
179 show an entirely different pattern; the leaping form (*C. guereza*) has both anterior and lateral canals
180 that are statistically indistinguishable from its congener, but possesses posterior canals that are
181 significantly smaller; $t = -2.656$, $df = 10$, Sig. (2-tailed) = 0.024 (Fig. 5).

182

183 Figure 4 about here

184

185 Figure 5 about here

186

187 Discussion

188

189 In *Presbytis*, the difference (larger lateral canals) between species that vary widely in the amount
190 they leap during locomotion is in the same canal and in the same direction as that reported for other
191 primates (Spoor et al., 2007). Combined with the fact that this statistically significant difference is
192 found even at relatively small sample sizes ($n=4-6$), this could be construed as support for previous
193 interpretations. The fact that *Colobus* species show a completely different pattern (the leaping form
194 has smaller posterior canals), however, suggest that caution should be exercised when attempting to
195 use the *Presbytis* results on their own to infer behaviour in extinct forms.

196

197 The failure of primate congeners that differ in specifics in the locomotor repertoire to show a
198 consistent pattern in semicircular canal size suggests that either a) leaping reflects a sufficiently
199 different aspect of locomotor function from the multivariate agility categories utilised previously
200 that the morphological signals in the organ of balance are not comparable, b) increasing the amount
201 of leaping in the locomotor repertoire by over 100% has no effect on labyrinth morphology, or c)
202 the influence of drift at the lower taxonomic level examined here is greater than any inferred

functional adaptation. Precisely which of these possibilities is most probable is unclear at present.

Even though it had been suggested previously (Spoor et al., 2007; Walker et al., 2008) that leaping may have a critical effect on the organ of balance, the differences between the broad behavioural categories of agility used in more wide-ranging studies on the one hand and specifically leaping on the other may be sufficient to result in distinct patterns of labyrinthine morphology. As the agility scores used in previous research (e.g., Spoor et al., 2007) include speed (and, presumably, additional attributes of locomotor behaviour), it may be that these factors (rather than percentage of leaping) have affected the results. Indeed, although the larger lateral canals of tarsiers have been linked explicitly to their leaping behaviour (Spoor et al., 2007), they have also been suggested to be the result of the increased degree of head rotation associated with their vertical clinging and leaping mode of movement (Spoor and Zonneveld, 1998). If this is the case, leaping may not be a sufficiently powerful selective pressure to affect the organ of balance substantially, in which case other explanations for the significant results obtained above are necessary.

It also could be argued that the amount of difference between taxa in terms of the percentage of leaping in the locomotor repertoire recorded here (over 100% in both cases) is insufficient to affect the organ of balance. If so, then the labyrinth differs from other anatomical systems (e.g., skeletal and muscular; Fleagle, 1976, 1977) that demonstrate just such variation within the same subfamily; indeed, for one case, in the same genus. Again, however, the contradictory nature of the results complicates the interpretation, as the significant results suggest that some effect is present, but in two different directions on two different canals. It has also been demonstrated that a substantial amount of phylogenetic signal may be present in labyrinthine morphology of some primate groups (Lebrun et al., 2010), which could affect the results, although the use of congeneric pairs reduces the likelihood that this is a significant factor. In addition, because behavioural flexibility is a characteristic of primates, and because behaviour can change more quickly than morphology,

229 evolutionary lag might be invoked (Blomberg et al., 2003), although it would not fully account for
230 the conflicting significant differences.

231

232 A final potential interpretation is that the differences seen are the result of genetic drift. If this is the
233 case, it would imply that chance affects labyrinth morphology to a greater extent than forces
234 produced during locomotion. In some ways, drift is (and perhaps should be) the null explanation
235 for differences between closely related taxa (or, indeed, individuals); a notion that is receiving more
236 attention recently (e.g., Weaver et al., 2007; Betti et al., 2010). Additional testing would be required
237 to see if the pattern of the variation in semicircular canal radius in colobines is distinguishable from
238 those derived from drift models; if not, no functional explanation for the observed differences
239 would be necessary. Alternatively, it may be that a multivariate approach (Gunz et al., 2012) is
240 needed to differentiate between taxa such as those investigated here, although this may limit the
241 extent to which individual traits of the labyrinth can be subjected to statistical testing. In any event,
242 the fact that these significant patterns are diametrically opposed in two sets of closely related taxa
243 reduces the confidence that can be placed on inferences derived from comparisons between fossil
244 and living taxa.

245

246 ACKNOWLEDGMENTS

247 The authors acknowledge financial support provided via a Primate Society of Great Britain grant
248 awarded to PMJ and a Japan Society for the Promotion of Science Fellowship awarded to TCR.
249 Permission to study the NHM material was facilitated by Roberto Portela Miguez. Scanning at
250 HYMS was generously provided by Paul O'Higgins and Michael J Fagan, and performed by Sue
251 Taft. The manuscript was improved by comments from the Editor and an anonymous reviewer. .

252

253 REFERENCES

254

255 Betti L, Balloux F, Hanihara T, Manica A (2010). The relative role of drift and selection in shaping
 256 the human skull. *American Journal of Physical Anthropology* 141:76–82.
 257

258 Billet G, Hautier L, Asher RJ, Schwarz C, Crumpton N, Martin T, Ruf I (2012). High morphological
 259 variation of vestibular system accompanies slow and infrequent locomotion in three-toed sloths.
 260 *Proceedings of the Royal Society B* 279: 3932-3939.
 261

262 Birkhead TR, Clarkson K, Reynolds MD, Koenig WD (1992). Copulation and mate guarding in the
 263 yellow-billed magpie *Pica nuttalli* and a comparison with the black-billed magpie *P. pica*.
 264 *Behaviour* 121: 110-130.
 265

266 Blomberg SP, Garland T, Ives AR (2003). Testing for phylogenetic signal in comparative data:
 267 behavioral traits are more labile. *Evolution* 57: 717-745.
 268

269 Boistel R, Herrel A, Lebrun R, Daghfous G, Tafforeau P, Losos JB, Vanhooydonck B (2011). Shake
 270 rattle and roll: the bony labyrinth and aerial descent in squamates. *Integrative and Comparative*
 271 *Biology* 51: 957–968.
 272

273 Crews D (1997). Species diversity and the evolution of behavioural controlling mechanisms. *Annals*
 274 *of the New York Academy of Sciences* 807: 1-21.
 275

276 Davison GWH (1982). Convergence with terrestrial cercopithecines by the monkey *Rhinopithecus*
 277 *roxellanae*. *Folia Primatologica* 37: 209-215.
 278

279 Dunham NT, Kane EE, McGraw WS (2015). Scapular morphology and forelimb use during
 280 foraging in four sympatric cercopithecids. *Folia Primatologica* 86: 474-489.

281

282 Ekdale EG (2013). Comparative anatomy of the bony labyrinth (inner ear) of placental mammals.
283 *PLoS ONE* 8(6): e66624. doi:10.1371/journal.pone.0066624

284

285 Fleagle JG (1976). Locomotor behaviour and skeletal anatomy of sympatric Malaysian leaf -
286 monkeys (*Presbytis obscura* and *Presbytis melalophos*). *Yearbook of Physical Anthropology* 20:
287 440-453.

288

289 Fleagle JG (1977). Locomotor behavior and muscular anatomy of sympatric Malaysian leaf -
290 monkeys (*Presbytis obscura* and *Presbytis melalophos*). *American Journal of Physical*
291 *Anthropology* 46: 297-308.

292

293 Fleagle JG (1999). *Primate adaptation and evolution*. 2nd ed. San Diego, CA, Academic Press

294

295 Fuentes A (1996). Feeding and ranging in the Mentawai Island langur (*Presbytis potenziani*).
296 *International Journal of Primatology* 17: 525-548.

297

298 Garland T, Adolph S. (1994). Why not to do two-species comparative studies: Limitations on
299 inferring adaptation. *Physiological Zoology* 67: 797-828.

300

301 Gebo DL, Chapman CA (1995). Positional behaviour in five sympatric Old World monkeys.
302 *American Journal of Physical Anthropology* 97: 49-76.

303

304 Gibson JJ (1968). *The senses considered as perceptual systems*. London, George Allen and Unwin.

305

306 Gray AA (1908). An investigation on the anatomical structure and relationships of the labyrinth in

307 the reptile, the bird, and the mammal. *Proceedings of the Royal Society B* 80: 507-528.

308

309 Grohé C, Tseng J, Lebrun R, Boistel R, Flynn JJ (2016). Bony labyrinth shape variation in extant

310 Carnivora: a case study of Musteloidea. *Journal of Anatomy* 228: 366–383

311

312 Gunz P, Ramsier M, Kuhrig M, Hublin J-J, Spoor F (2012). The mammalian bony labyrinth

313 reconsidered, introducing a comprehensive geometric morphometric approach. *Journal of Anatomy*

314 220: 529-543.

315

316 Johnson PM (2012). *Leaping in the ears? A test of the semicircular canals as locomotor indicators*

317 *in closely related primate taxa*. MRes thesis, University of Roehampton.

318

319 Katz PS, Harris-Warrick RM (1999). The evolution of neuronal circuits underlying species-specific

320 behaviour. *Current Opinion in Neurobiology* 9: 628–633.

321

322 Lebrun R, de Leon MP, Tafforeau P, Zollikofer C (2010). Deep evolutionary roots of strepsirrhine

323 primate labyrinthine morphology. *Journal of Anatomy* 216:368–80.

324

325 Lieberman DE (2011). *The evolution of the human head*. Harvard, Harvard University Press.

326

327 Martin RD, Ross CF (2005). The evolutionary and ecological context of primate vision. In: *The*

328 *primate visual system: a comparative approach* (Kremers J, ed.), pp. 1-36. Chichester, John Wiley

329 and Sons.

330

331 Mather G (2009). *Foundations of sensation and perception*. Hove, Psychology Press.

332

333 Meyer D, Rinaldi ID, Ramlee H, Perwitasari-Farajallah D, Hodges JK, Roos C (2011).
 334 Mitochondrial phylogeny of leaf monkeys (genus *Presbytis*, Eschscholtz, 1821) with implications
 335 for taxonomy and conservation. *Molecular Phylogenetics and Evolution* 59: 311–319
 336
 337 McGraw WS (1998). Comparative locomotion and habitat use of six monkeys in the Tai Forest,
 338 Ivory Coast. *American Journal of Physical Anthropology* 105:493-510.
 339
 340 Pfaff C, Martin T, Ruf I (2015). Bony labyrinth morphometry indicates locomotor adaptations in the
 341 squirrel-related clade (Rodentia, Mammalia). *Proceedings of the Royal Society B* 282, Article
 342 Number: 20150744 .
 343
 344 Reed K (1999). Population density of primates in communities: Differences in community structure.
 345 In: *Primate communities* (Fleagle JG, Janson C, Reed KE, eds.), pp. 116-140. Cambridge,
 346 Cambridge University Press.
 347
 348 Rowe, N., 1996. The pictorial guide to the living primates. East Hampton, NY, Pogonias Press.
 349
 350 Ryan TM, Silcox MT, Walker A, Mao X, Begun DR, Benefit BR, Gingerich PD,
 351 Köhler M, Kordos L, McCrossin ML, Moyà -Solà S, Sanders WJ, Seiffert ER, Simons E, Zalmout
 352 IS, Spoor F (2012). Evolution of locomotion in Anthropoidea: The semicircular canal evidence.
 353 *Proceedings of the Royal Society B* 279: 3467-3475.
 354
 355 Schmelzle T, Maier W, Sánchez-Villagra MR (2007). Vestibular labyrinth evolution in
 356 diprotodontian marsupial mammals. *Mammal Study* 32:83-97.
 357
 358 Sinnatamby CS (2006). *Last's anatomy: regional and applied, 11th Ed.* Churchill Livingstone,

359 Edinburgh.

360

361 Spoor F (2003). The semicircular canal system and locomotor behaviour, with special reference to

362 hominin evolution. *Courier Forschungsinstitut Senckenberg* 243: 93-104.

363

364 Spoor F, Zonneveld F (1995). Morphometry of the primate bony labyrinth: A new method based on

365 high-resolution computed tomography. *Journal of Anatomy* 186: 271-286.

366

367 Spoor F, Zonneveld F (1998) Comparative review of the human bony labyrinth. *Yearbook of*

368 *Physical Anthropology* 27: 211-251.

369

370 Spoor F, Garland T Jr, Krovitz G, Ryan TM, Silcox MT, Walker A (2007). The primate semicircular

371 canal system and locomotion. *Proceedings of the National Academy of Sciences of the United States*

372 *of America* 104: 10808-10812.

373

374 Su DF, Jablonski NG (2009). Locomotor behavior and skeletal morphology of the odd-nosed

375 monkeys. *Folia Primatologica* 80: 189-219.

376

377 Ting N (2008). Mitochondrial relationships and divergence dates of the African colobines: evidence

378 of Miocene origins for the living colobus monkeys. *Journal of Human Evolution* 55: 312-325

379

380 Walker A, Ryan TM, Silcox MT, Simons E, Spoor F (2008). The semicircular canal system and

381 locomotion: The case of extinct lemuroids and lorisooids. *Evolutionary Anthropology* 17: 135-145.

382

383 Wang XP, Yu L, Roos C, Ting N, Chen CP, Wang J, Zhang YP (2012). Phylogenetic relationships
384 among the colobine monkeys revisited: new insights from analyses of complete mt genomes and 44
385 nuclear non-coding markers. *PLoS ONE* 7(4): e36274. doi:10.1371/journal.pone.0036274
386
387 Weaver T, Roseman C, Stringer C (2007). Were Neandertal and modern human cranial differences
388 produced by natural selection or genetic drift? *Journal of Human Evolution* 53: 135-145.
389
390 Welker KL, Orkin JD, Ryan TM (2009). Analysis of intraindividual and intraspecific variation in
391 semicircular canal dimensions using high-resolution x-ray computed tomography. *Journal of*
392 *Anatomy* 215: 444-451.

393 TABLES

394

395 Table 1: Percentage of leaping in the locomotor repertoire of *Presbytis* and *Colobus*. Data derived
396 from Fuentes (1996), Gebo and Chapman (1995), McGraw (1998) and Reed (1999). For both
397 genera, the species indicated by an asterisk (*) leaps more than twice as frequently as its congener.

398

399	TAXON	% leaping	% quadrupedalism	% other
400	-----			
401	<i>Presbytis melalophos</i> *	67.5	20.5	12
402	<i>P. potenziani</i>	25	75	0
403	<i>Colobus guereza</i> *	38	41	21
404	<i>C. polykomos</i>	14.5	71.2	14.3
405	-----			

406

407 Table 2. Species means for canal radius R (mm) and the cranial size proxy (prosthion-inion length;
408 mm)

409	-----					
410	Taxon	n	Anterior	Posterior	Lateral	Size
411	<i>Presbytis melalophos</i>	5	2.24	2.23	2.34	98.62
412	<i>Presbytis potenziani</i>	4	2.26	2.00	1.97	92.98
413	<i>Colobus guereza</i>	6	2.66	2.32	2.22	112.96
414	<i>Colobus polykomos</i>	6	2.61	2.48	2.28	109.28
415	-----					

416

417

418 FIGURE LEGENDS

419

420 Figure 1: Location of the bony labyrinth (shaded) in the temporal bone, seen in a 3D virtual
421 reconstruction of a *Presbytis* cranium derived from μ CT scans and rendered semi-transparent.

422

423 Figure 2: 3D virtual reconstruction of the bony labyrinth of *Presbytis*. The semicircular canals are
424 orientated orthogonally to one another for sensing pitch, roll and yaw of the head. Scale bar = 1mm.

425

426 Figure 3: Height (h) and width (w) measurements used to calculate canal radius (R). Scale bar =
427 1mm.


428

429 Figure 4: Virtual canals of a) *Colobus guereza*, b) *C. polykomos* (reversed), c) *Presbytis*
430 *melalophos*, and d) *P. potenziani*. Scale bars = 1mm.

431


432 Figure 5: Box and whiskers plots of size-adjusted R of *Presbytis* and *Colobus* semicircular canals.
433 The anterior, posterior and lateral canals are indicated below each chart. The species that leaps
434 more frequently is indicated by the monkey symbol.

435




436

437







438

439



440


441



442
443



 *Presbytis melalophos*  *P. potenziani*



 *Colobus guereza*  *C. polykomos*

